



LAWRENCE  
LIVERMORE  
NATIONAL  
LABORATORY

# Comparing the gain of the Ne K- $\alpha$ inner-shell X-ray laser using the X-FEL to drive the kinetics with photo-ionization versus photo-excitation

J. Nilsen

May 20, 2014

International Conference on X-Ray Lasers 2014  
Fort Collins, CO, United States  
May 25, 2014 through May 30, 2014

## **Disclaimer**

---

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

# *Comparing the gain of the Ne K- $\alpha$ inner-shell X-ray laser using the X-FEL to drive the kinetics with photo-ionization versus photo-excitation*

May 27, 2014

presented at 14<sup>th</sup> International Conference on X-ray Lasers

Fort Collins, Colorado, 25 – 30 May, 2014

Joseph Nilsen

*Lawrence Livermore National Laboratory*



LLNL-CONF-654625

This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344. Lawrence Livermore National Security, LLC




## Can resonant photo-excitation allow us to produce inner shell X-ray lasers more efficiently?

- Review the photoionization pumped inner-shell Ne X-ray laser
- Describe Ne model that includes photo-excitation processes
- Model an inner-shell Ne XRL pumped by photo-excitation
- Examine sensitivity of Ne XRL gain to XFEL linewidth and duration
- Discuss future possibilities for X-ray lasers driven by X-FEL' s

# Photoionization of a 1s electron in neutral Ne I creates a highly excited state of singly ionized Ne II that lases on the K- $\alpha$ line at 848.6 eV as demonstrated at LCLS-XFEL

P.I. cross section is 0.3 Mb


Auger lifetime is 2.3 fsec



## LCLS XFEL can be used to longitudinally pump an X-ray laser plasma and replace a traditional line source

Consider an XFEL beam with  $10^{12}$  photons in 100 fs pulse with 0.1% bandwidth focused to 1  $\mu\text{m}$  spot as the baseline drive



Gives flux of  $10^{33}$  photons/[sec  $\text{cm}^2$ ]



## Photoionization of 1s electron in Ne I with 875 eV XFEL can create large gain on K- $\alpha$ line of Ne II at 848.6 eV that can be increased by reducing time duration of XFEL


Baseline XFEL beam has  $10^{12}$  photons in 0.9 eV linewidth focused to 1  $\mu\text{m}$  spot  
Figures look at sensitivity to reducing number of photons by multiplying photon flux by number from 1 to 0.001

Time is relative to peak of XFEL pulse: consider 100 fs vs 1 fs FWHM XFEL pulses



# Photo-excitation of the 1s-3p transition in neutral Ne I creates a highly excited state of neutral Ne I that lases on the K- $\alpha$ lines at 848.96 eV

P.E. cross section is 18 Mb





## Photo-excitation of the 1s-3p transition in neutral Ne I could pump a Ne XRL at 849 eV using much less energy than the photoionization driven laser

Consider an XFEL beam with  $10^{12}$  photons in 100 fs pulse with 0.1% bandwidth focused to 1  $\mu\text{m}$  spot as the baseline drive

Gives flux of  $10^{33}$  photons/[sec  $\text{cm}^2$ ]

Photoionization rate is  $3.0 \times 10^{14}$  / sec for XFEL at 875 eV

Photo-excitation rate is  $7.5 \times 10^{15}$  / sec for XFEL tuned to 1s-3p

Experiments had  $L=0.28$  cm and  $N_{\text{ion}}=2 \times 10^{19}/\text{cc}$

Advantages of photoionization:

1. Not resonant so all of energy used to drive photoionization
2. Insensitive to bandwidth of XFEL

Advantage of photo-excitation:

1. Much higher excitation rate than photoionization



Disadvantage of photo-excitation:

1. Need to tune XFEL to be resonant
2. Require narrow bandwidth matched to linewidth of transition
3. Higher opacity on line being pumped

## Photo-excitation of 1s-3p line in Ne I with 867.6 eV XFEL can create large gain on K- $\alpha$ line of Ne I at 849 eV that can be increased by reducing time duration of XFEL



Baseline XFEL beam has  $10^{12}$  photons in 0.9 eV linewidth focused to 1  $\mu\text{m}$  spot  
Figures look at sensitivity to reducing number of photons by multiplying photon flux by  
number from 1 to 0.001

Time is relative to peak of XFEL pulse: consider 100 fs vs 1 fs FWHM XFEL pulses




## Photo-excitation of 1s-3p line in Ne I with 867.6 eV XFEL can create larger gain at much lower flux than using photoionization of 1s electron in Ne I at 875 eV

Baseline XFEL beam has  $10^{12}$  photons in 0.9 eV linewidth focused to 1  $\mu\text{m}$  spot  
Figures look at sensitivity to reducing number of photons by multiplying photon flux by number from 1 to 0.001




**Photoionization of a 2p or 2s electron in neutral Ar I creates a highly excited state of singly ionized Ar II that could lase on the 3s-2p or 3p-2s lines near 233 and 311 eV**



# Photoionization of a 2p electron in Cu I (or Cu II) creates a highly excited state of Cu II (or Cu III) that could lase on the 3d-2p line near 922 eV

Photo-excitation of 2p to 4d level would also drive 3d-2p laser



## Resonant photo-excitation could allow us to produce inner shell X-ray lasers more efficiently

- Reviewed the photoionization pumped inner-shell Ne X-ray laser
- Described Ne model that includes photo-excitation processes
- Model an inner-shell Ne XRL pumped by resonant photo-excitation
- Reducing the pulse duration from 100 fs to 1 fs increases the gain significantly for both the photoionization and photo-excitation driven inner shell Ne laser
- Driving the inner shell Ne laser with resonant photo-excitation can reduce the flux requirements by several orders of magnitude
- Reducing the linewidth of the XFEL to match the linewidth of the resonant photo-excitation transition should further reduce the flux requirements
- New inner shell schemes that lase on 3-2 transitions using L-shell photoionization look promising for future